


**Prior Authorization Review Panel  
MCO Policy Submission**

A separate copy of this form must accompany each policy submitted for review.  
Policies submitted without this form will not be considered for review.

<b>Plan: AmeriHealth Caritas Pennsylvania Community Health Choices</b>	<b>Submission Date: 5/1/2024</b>
<b>Policy Number: ccp.1455</b>	<b>Effective Date: 5/2020</b> <b>Revision Date: April 1, 2024</b>
<b>Policy Name: Stereotactic navigation in sinus surgery</b>	
<b>Type of Submission – Check all that apply:</b>  <input checked="" type="checkbox"/> <b>New Policy</b> <input type="checkbox"/> <b>Revised Policy*</b> <input type="checkbox"/> <b>Annual Review – No Revisions</b> <input type="checkbox"/> <b>Statewide PDL</b>	
<b>*All revisions to the policy <u>must</u> be highlighted using track changes throughout the document.</b>  <b>Please provide any clarifying information for the policy below:</b>  <b>See tracked changes below.</b>	
<b>Name of Authorized Individual (Please type or print):</b>  Manni Sethi, MD, MBA, CHCQM	<b>Signature of Authorized Individual:</b>  

# Stereotactic navigation in sinus surgery

Clinical Policy ID: CCP.1455

Recent review date: 4/2024

Next review date: 8/2025

Policy contains: Computer-assisted sinus surgery; endoscopic sinus surgery; image-guided sinus surgery; surgical navigation.

*AmeriHealth Caritas Pennsylvania Community HealthChoices has developed clinical policies to assist with making coverage determinations. AmeriHealth Caritas Pennsylvania Community HealthChoices' clinical policies are based on guidelines from established industry sources, such as the Centers for Medicare & Medicaid Services (CMS), state regulatory agencies, the American Medical Association (AMA), medical specialty professional societies, and peer-reviewed professional literature. These clinical policies along with other sources, such as plan benefits and state and federal laws and regulatory requirements, including any state- or plan-specific definition of "medically necessary," and the specific facts of the particular situation are considered by AmeriHealth Caritas Pennsylvania Community HealthChoices when making coverage determinations. In the event of conflict between this clinical policy and plan benefits and/or state or federal laws and/or regulatory requirements, the plan benefits and/or state and federal laws and/or regulatory requirements shall control. AmeriHealth Caritas Pennsylvania Community HealthChoices' clinical policies are for informational purposes only and not intended as medical advice or to direct treatment. Physicians and other health care providers are solely responsible for the treatment decisions for their patients. AmeriHealth Caritas Pennsylvania Community HealthChoices' clinical policies are reflective of evidence-based medicine at the time of review. As medical science evolves, AmeriHealth Caritas Pennsylvania Community HealthChoices will update its clinical policies as necessary. AmeriHealth Caritas Pennsylvania Community HealthChoices' clinical policies are not guarantees of payment.*

## Coverage policy

Stereotactic navigation (also known as three-dimensional image guidance or computer-aided surgery) is considered standard of care when added to general endoscopic sinus surgery; therefore, it is not separately reimbursable as an add-on to the general procedure.

### Limitations

No limitations were identified during the writing of this policy.

### Alternative covered services

- Clinical examination.
- Diagnostic rhinoscopy.
- Endoscopic or open sinus surgery.
- Preoperative and postoperative radiography, computed tomography, or magnetic resonance imaging.
- Professional consultation.

## Background

Sinus surgery is one of the most frequently performed surgical operations in the United States, and the most common indication is chronic refractory rhinosinusitis. Since the late 1990s, there has been a shift from open procedures to endoscopic procedures and a decrease in the incidence of major complications associated with

endoscopic approaches in adults. For example, in England from 2010 to 2019, the number of endoscopic sinus procedures increased by 21.1%, and the number of open procedures declined by 35.3% (Gupta, 2020).

Rhinosinusitis is a clinical diagnosis, dependent on clinical and rhinoscopic findings. Some diseases involving the sinonasal region and skull base are inaccessible to sinonasal endoscopy or are not directly visible on clinical examination. In such cases, imaging can determine the extent and possible nature of the disease and provide guidance for surgical planning.

Multidetector or cone-beam computed tomography is the preferred imaging modality for surgical planning in most circumstances for its clear delineation of the complex ethmoidal anatomy, ostiomeatal unit, and anatomic variations, including the presence of sphenoidal (Onodi) air cells (American College of Radiology, 2021). Magnetic resonance imaging of the face and sinuses with inclusion of adjacent brain and orbits is preferred for soft-tissue masses, delineation of orbital, skull base, or intracranial involvement, and for suspected invasive fungal sinusitis.

Endoscopic sinus surgery is a set of minimally invasive techniques that apply endoscopic instrumentation to directly examine the sinuses, remove sinus blockages, and restore sinus ventilation and normal function. It is a treatment of choice for chronic rhinosinusitis refractory to medical treatment, and its role is expanding in the management of other sinus, orbit, and skull base diseases (Homsy, 2022).

Endoscopic sinus surgery is a technically demanding procedure due to anatomical complexity, variation in normal anatomy, and proximity to critical structures such as the olfactory fossa, skull base, vascular structures, and orbit. Anatomic distortion from chronically inflamed mucosa and absent anatomic landmarks from prior surgery further increase both potential risk of major and minor complications and operating time.

In a systematic review/meta-analysis of 13 studies (n = 1,274) of complications from endoscopic sinus surgery for chronic rhinosinusitis, the complication rate was 5.3% with enhanced recovery after surgery – a multimodal multi-disciplinary planning program – compared to the 19.1% rate using standard care (Wu, 2023). To optimize the visual surgical field and mitigate the risk of complications, stereotactic navigation has been proposed as an adjunct in endoscopic and external-approach sinus surgery.

The image guidance systems consist of a computer workstation, tracking system, and specially designed navigation instruments (Oakley, 2016). They require a compatible preoperative computed tomography scan of the sinuses using a special image guidance protocol, for example, a non-contrast axial image with 1 millimeter or thinner cuts. Image guidance systems digitally link the images and display them in axial, coronal, and sagittal planes. The image is coupled to the surgical instruments by an electromagnetic or optical tracking system. This gives the surgeon the ability to navigate the surgical instruments in real time within the diseased sinuses and improve the spatial orientation with vital structures while watching the monitor.

## Findings

For this policy, we included two systematic reviews (Ramakrishnan, 2013; von Hofsten, 2013), two meta-analyses (Dalgorf, 2013; Vreugdenburg, 2016), and one consensus-based professional guideline (American Academy of Otolaryngology — Head and Neck Surgery, 2023) to address whether the perceived net benefit of stereotactic guidance translates into a clinically meaningful benefit and to identify the optimal candidates for the procedure. The evidence from the primary studies included in the systematic reviews and meta-analyses was of low-to-moderate quality and comprised of retrospective cohorts and case series of adult and pediatric patients from tertiary care settings. In general, allocation of stereotactic navigation was based on availability of equipment with likely selection bias toward those with more severe disease.

The perceived potential of increased safety and improved surgical outcomes with stereotactic navigation in endoscopic sinus surgery led to the introduction and diffusion of the technology in the early 2000s in the absence

of a clear benefit demonstrated in clinical research. A survey of members of the American Rhinologic Society showed a growth in both the availability of stereotactic navigation to most practitioners and its use from 2005 to 2010 ( $P < .0001$ ) (Justice, 2012). Since stereotactic navigation is widely available, randomized controlled studies would be impractical to perform at this point. Moreover, major complications in endoscopic sinus surgery are relatively rare, and sufficiently powered studies needed to detect statistically meaningful differences in outcomes attributable to stereotactic navigation would be prohibitive.

To illustrate, two qualitative systematic reviews (Ramakrishnan, 2013; von Hofsten, 2013) failed to show a definitive reduction in complications or definitive improvement in outcomes. Their conclusions may reflect the inability to carry out more definitive studies rather than a lack of efficacy, per se. The individual primary studies lacked the power to identify a statistically meaningful effect of stereotactic navigation added to endoscopic sinus surgery, if one existed.

Pooling results through meta-analysis from multiple studies may increase the statistical power over individual studies to detect a meaningful effect. A meta-analysis (Dalgorf, 2013) of 14 comparative cohorts found both major complications ( $P = .007$ ) and total complications ( $P = .02$ ) were more common in the non-image guided group, but there were no significant between-group differences in risk of orbital complications, intracranial complications, major hemorrhage, completion of operation, revision surgery, or patient-reported outcome measures.

Another meta-analysis (Vreugdenburg, 2016) of nine studies demonstrated a statistically significant decrease in the likelihood of total, major, and orbital complications in participants who had stereotactic navigation. However, a statistical benefit in reducing surgical revisions, major hemorrhage, or minor complications was not observed.

The retrospective nature of the primary studies complicates the ability to identify the optimal candidate for stereotactic guidance, along with procedure complexity, variation in patient anatomy, inflammatory or neoplastic disease burden, and surgeon skill and training (Dalgorf, 2013). American Rhinologic Society survey respondents identified primary anterior ethmoidectomy, revision anterior ethmoidectomy, primary total ethmoidectomy, Lothrop procedure, cerebrospinal fluid leak repair, tumor surgery, orbital decompression, and optic nerve decompression (all  $P < .05$ ) as major indications for stereotactic navigation (Justice, 2012). These indications illustrate that stereotactic navigation is best reserved for situations for which major surgical complications would be the greatest or in which suboptimal or incomplete surgery may lead to poorer outcomes or revision surgery.

The American Academy of Otolaryngology — Head and Neck Surgery (2023) recommends indications for stereotactic navigation based on consensus and current evidence and stresses it would be impossible to corroborate with Level 1 evidence. The technology should be used at the discretion of the operating surgeon in clarifying complex anatomy during sinus and skull base surgery for the following indications:

- Revision sinus surgery.
- Distorted sinus anatomy of development, postoperative, or traumatic origin.
- Extensive sino-nasal polyposis.
- Pathology involving the frontal, posterior ethmoid, or sphenoid sinuses.
- Disease abutting the skull base, orbit, optic nerve, or carotid artery.
- Cerebrospinal fluid rhinorrhea or conditions where there is a skull base defect.
- Benign and malignant sino-nasal neoplasms.

In 2021, we added one cost-effectiveness study based on the results of pediatric patients who underwent sinus surgery with intraoperative navigation ( $n = 60$ ) and without intraoperative navigation ( $n = 59$ ) between 2003 and 2016. Use of intraoperative navigation was associated with more complex surgeries with more sinuses opened ( $P = .008$ ), increased presence of the attending surgeon and a trainee ( $P < .001$  for both), and longer surgical time ( $P < .001$ ). Ultimately, intraoperative navigation did not decrease complication rates or revision rates and

was not cost-effective. It was used primarily as an educational tool or to increase confidence in intraoperative identification of landmarks. These results warrant no policy changes.

In 2022, we updated the American Academy of Otolaryngology – Head and Neck Surgery position statement and added a guideline from the International Stereotactic Radiosurgery Society that recommended stereotactic radiosurgery as a treatment for grade 1 meningioma (Marchetti, 2020). We also added several large reviews:

- A systematic review/meta-analysis of 27 studies found that stereotactic radiosurgery for benign intracranial meningiomas had a local control rate from 71% to 100% and progression-free-survival rate from 55% to 97% after 10 years, with a low toxicity rate (Marchetti, 2020).
- A systematic review of 47 studies on endoscopic sinus surgery found the most commonly identified significant landmarks were maxillary sinus ostium; orbital wall; frontal recess; skull base; ground lamella; fovea posterior; and sphenoid sinus ostium. These landmarks may lead to systematic practice guidelines in navigated endoscopic sinus surgery (Baudoin, 2021).

In 2023, we added two systematic review/meta-analyses of:

- Twenty-one studies (n = 705) of patients treated with stereotactic radiosurgery for dural arteriovenous fistulas. Non-cavernous sinus fistulas were associated with lower rates of complete obliteration ( $P = .03$ ) and symptom cure rate ( $P = .001$ ) (Singh, 2022).
- Five studies (n = 208) of patients with pituitary adenomas who underwent endoscopic resection of the medial wall of the cavernous sinus showed the prevalence of medial wall invasion varied from 10.4% to 36.7%. Relative success of various techniques is discussed (de Macedo Filho, 2022).

In 2024, we updated the references and added a randomized study (n = 200) of participants undergoing endoscopic sinus surgery with versus without an image-guided system in China. Four re-hospitalizations and re-operation due to bleeding for the group without image guidance occurred, compared with none for the image guidance group. Total costs were slightly greater for the group without image guidance, due to re-hospitalizations (Wang, 2023).

We also added a meta-analysis of 21 studies (n = 199) of participants undergoing surgery for cavernous sinus hemangiomas. One-hundred forty-nine of the participants received standalone stereotactic radiosurgery, of whom 72.4% had greater than a 50% decrease in tumor volume, after being followed for an average of three years (Mishra, 2023).

## References

On January 25, 2024, we searched PubMed and the databases of the Cochrane Library, the U.K. National Health Services Centre for Reviews and Dissemination, the Agency for Healthcare Research and Quality, and the Centers for Medicare & Medicaid Services. Search terms were “otorhinolaryngologic surgical procedures (MeSH),” “paranasal sinus diseases/surgery (MeSH),” and “Surgery, Computer-Assisted (MeSH),” and “endoscopic sinus surgery.” We included the best available evidence according to established evidence hierarchies (typically systematic reviews, meta-analyses, and full economic analyses, where available) and professional guidelines based on such evidence and clinical expertise.

American Academy of Otolaryngology — Head and Neck Surgery. Position statement: Intra-operative use of computer aided surgery. <https://www.entnet.org/resource/position-statement-intra-operative-use-of-computer-aided-surgery/#:~:text=Position%20Statement%3A%20Intra%2DOperative%20Use%20of%20Computer%20Aided%20Surgery.->

[Position Statement 3A Intra&text=The American Academy of Otolaryngology, sinus and skull base surgery](#). Revised October 3, 2023.

American College of Radiology. ACR Appropriateness Criteria® sinonasal disease. <https://acsearch.acr.org/docs/69502/Narrative/>. Revised 2021.

Baudoin T, Greguric T, Bacan F, Jelavic B, Geber G, Kosec A. A systematic review of common landmarks in navigated endoscopic sinus surgery (NESS). *Comput Assist Surg (Abingdon)*. 2021;26(1):77-84. Doi: 10.1080/24699322.2021.1992504.

Dalgorf DM, Sacks R, Wormald P-J, et al. Image-guided surgery influences perioperative morbidity from endoscopic sinus surgery: A systematic review and meta-analysis. *Otolaryngol Head Neck Surg*. 2013;149(1):17-29. Doi: 10.1177/0194599813488519.

De Macedo Filho LJM, Diogenes AVG, Barreto EG, et al. Endoscopic endonasal resection of the medial wall of the cavernous sinus and its impact on outcomes of pituitary surgery: A systematic review and meta-analysis. *Brain Sci*. 2022;12(10):1354. Doi: 10.3390/brainsci12101354.

Gupta KK, Jolly K, Bhamra N, Osborne MS, Ahmed SK. The evolution of sinus surgery in England in the last decade – An observational study. *World J Otorhinolaryngol Head Neck Surg*. 2020;7(3):240-246. Doi: 10.1016/j.wjorl.2020.10.002.

Homsy MT, Gaffey MM. Sinus endoscopic surgery. In: *StatPearls* [Internet]. Treasure Island (FL): Stat Pearls Publishing; 2024 Jan. <https://www.ncbi.nlm.nih.gov/books/NBK563202>. Last updated September 12, 2022. Justice JM, Orlandi RR. An update on attitudes and use of image-guided surgery. *Int Forum Allergy Rhinol*. 2012;2(2):155-159. Doi: 10.1002/alr.20107.

Marchetti M, Sahgal A, De Salles AA, et al. Stereotactic radiosurgery for intracranial noncavernous sinus benign meningioma: International Stereotactic Radiosurgery Society systematic review, meta-analysis and practice guidelines. *Neurosurgery*. 2020;87(5):879-890. Doi: 10.1093/neuros/nyaa169.

Mishra S, Kumar AG, Garg K, et al. Role of stereotactic radiosurgery for cavernous sinus hemangiomas – An individual patient data-based meta-analysis. *Neurol India*. 2023;71(Supplement):S21-S30. Doi: 10.4103/0028-3886.373654.

Oakley GM, Barham HP, Harvey RJ. Utility of image-guidance in frontal sinus surgery. *Otolaryngol Clin North Am*. 2016;49(4):975-988. Doi: 10.1016/j.otc.2016.03.021.

Ramakrishnan VR, Orlandi RR, Citardi MJ, et al. The use of image-guided surgery in endoscopic sinus surgery: An evidence-based review with recommendations. *Int Forum Allergy Rhinol*. 2013;3(3):236-241. Doi: 10.1002/alr.21094.

Singh R, Chen C-J, Didwania P, et al. Stereotactic radiosurgery for dural arteriovenous fistulas: A systematic review and meta-analysis and International Stereotactic Radiosurgery Society practice guidelines. *Neurosurgery*. 2022;91(1):43-58. Doi: 10.1227/neu.0000000000001953.

Von Hofsten P, Bark S, Blank S, et al. Navigationsutrustning som stöd vid endoskopisk sinuskirurgi. [Image-guided system for endoscopic sinus surgery]. Region Västra Götaland, Sahlgrenska University Hospital, HTA-centrum. Report number 2013:60.

[https://www.researchgate.net/publication/273761926\\_Navigationsutrustning\\_som\\_stod\\_vid\\_endoskopisk\\_sinuskirurgi\\_Image-guided\\_system\\_for\\_endoscopic\\_sinus\\_surgery](https://www.researchgate.net/publication/273761926_Navigationsutrustning_som_stod_vid_endoskopisk_sinuskirurgi_Image-guided_system_for_endoscopic_sinus_surgery). Published 2013.

Vreugdenburg TD, Lambert RS, Atukorale YN, Cameron AL. Stereotactic anatomical localization in complex sinus surgery: A systematic review and meta-analysis. *Laryngoscope*. 2016;126(1):51-59. Doi: 10.1002/lary.25323.

Wang Z, Liu C, Tan B, et al. Clinical and economic benefits of image-guided system in functional endoscopic sinus surgery: A retrospective chart review study in China. *Cost Eff Resour Alloc.* 2023;21(1):1. Doi: 10.1186/s12962-023-00414-2.

Wu Y, Fu Y, He Y, et al. The application of enhanced recovery after surgery (ERAS) in chronic rhinosinusitis patients undergoing endoscopic sinus surgery: A systematic review and meta-analysis. *PLoS One.* 2023;18(9):e0291835. Doi: 10.1371/journal.pone.0291835.

## Policy updates

4/2020: initial review date and clinical policy effective date: 5/2020

4/2021: Policy references updated.

4/2022: Policy references updated.

4/2023: Policy references updated.

4/2024: Policy references updated.