


**Prior Authorization Review Panel
MCO Policy Submission**

A separate copy of this form must accompany each policy submitted for review.
Policies submitted without this form will not be considered for review.

Plan: AmeriHealth Caritas Pennsylvania	Submission Date: March 29, 2022
Policy Number: CCP.1448	Effective Date: 5/2020 Revision Date: March 1, 2022
Policy Name: Flexitouch® pneumatic compression devices for lymphedema after head and neck surgery	
Type of Submission – Check all that apply: <input type="checkbox"/> New Policy <input checked="" type="checkbox"/> Revised Policy* <input type="checkbox"/> Annual Review – No Revisions <input type="checkbox"/> Statewide PDL	
*All revisions to the policy <u>must</u> be highlighted using track changes throughout the document. Please provide any clarifying information for the policy below: Please see revisions below using tracked changes.	
Name of Authorized Individual (Please type or print): Akintayo Akinlawon, MD	Signature of Authorized Individual: 

Flexitouch® pneumatic compression devices for lymphedema after head and neck surgery

Clinical Policy ID: CCP.1448

Recent review date: 3/2022

Next review date: 7/2023

Policy contains: Flexitouch; head and neck cancer; lymphedema; pneumatic compression devices.

AmeriHealth Caritas Pennsylvania Community HealthChoices has developed clinical policies to assist with making coverage determinations. AmeriHealth Caritas Pennsylvania Community HealthChoices' clinical policies are based on guidelines from established industry sources, such as the Centers for Medicare & Medicaid Services (CMS), state regulatory agencies, the American Medical Association (AMA), medical specialty professional societies, and peer-reviewed professional literature. These clinical policies along with other sources, such as plan benefits and state and federal laws and regulatory requirements, including any state- or plan-specific definition of "medically necessary," and the specific facts of the particular situation are considered by AmeriHealth Caritas Pennsylvania Community HealthChoices when making coverage determinations. In the event of conflict between this clinical policy and plan benefits and/or state or federal laws and/or regulatory requirements, the plan benefits and/or state and federal laws and/or regulatory requirements shall control. AmeriHealth Caritas Pennsylvania Community HealthChoices' clinical policies are for informational purposes only and not intended as medical advice or to direct treatment. Physicians and other health care providers are solely responsible for the treatment decisions for their patients. AmeriHealth Caritas Pennsylvania Community HealthChoices' clinical policies are reflective of evidence-based medicine at the time of review. As medical science evolves, AmeriHealth Caritas Pennsylvania Community HealthChoices will update its clinical policies as necessary. AmeriHealth Caritas Pennsylvania Community HealthChoices' clinical policies are not guarantees of payment.

Coverage policy

Flexitouch® pneumatic compression devices (Tactile Systems Technology, Inc., Northbrook, IL) for lymphedema after head and neck surgery are investigational/not clinically proven and, therefore, not medically necessary.

Requests for this service will be evaluated on a case by case basis.

Limitations

No limitations were identified during the writing of this policy.

Alternative covered services

No alternative covered services were identified during the writing of this policy.

Background

Head and neck cancers are predominately squamous cell carcinomas and account for about 4% of all cancers in the United States (National Cancer Institute, 2021a). The majority of these cancers occur in men and in persons over the age of 50. Most head and neck cancers affect the epiglottis, esophagus, glottis, hard palate, hypopharynx, oral cavity, oropharynx, palatine tonsil, pharynx, subglottis, supraglottis, tongue, or trachea.

The majority of patients present with regional spread or distant metastasis at the time of diagnosis, as only approximately 28% of head and neck cancers are diagnosed when the cancer is confined to the primary site

(National Cancer Institute, 2021b). Estimated five-year survival rates after diagnosis for head and neck cancers range from 18% for cancers with distant metastasis to 85% for cancers confined to the primary site. While treatments can vary by cancer location, cancer stage, patient age, and general health, they may include chemotherapy, radiation therapy, surgery, and targeted chemotherapy (National Cancer Institute, 2021a).

Lymphedema is common in patients who have been treated for head and neck cancer, especially from surgery and radiation, as up to 75% of patients will manifest some signs and symptoms of lymphedema after treatment (Miller, 2016). Secondary lymphedema from head and neck cancer treatment affect the face, mouth, and neck, both internally and externally. It may manifest as limitations to communication (speaking, reading, writing, and hearing), alimentation, and respiration. In severe cases, vision may be impaired, and thus ambulation may be impeded. Psychological effects from physical disfiguration can also occur (Smith, 2010). A study of 103 patients with head and neck cancer found that lymphedema severity correlates with symptom burden, functional status, and quality of life (Deng, 2013).

The American Head and Neck Society states that the most commonly used treatment for lymphedema is complete decongestive therapy, which includes massage to drain lymph, compression bandages or clothing, exercises to improve lymph flow, and skin care (Miller, 2016). Most of these treatments can be performed at home, guided by a lymphedema therapist. Success rates are relatively high, especially for those using these treatments at least five times weekly over a three-month period. A study of 1,202 patients with head and neck cancer-related lymphedema showed 60% improved after complete decongestive therapy, and treatment adherence was a significant predictor ($P < .001$) of response (Smith, 2015).

A recently developed treatment for lymphedema after cancer treatment is the pneumatic compression device, a two-phase lymph preparation and drainage device that stimulates the lymphatic system to move lymph fluids from areas of impaired lymphatic function to healthy areas. Garments are made with stretch material that wraps around the affected area, so no fitting process is needed. They apply sequential inflation and deflation to several chambers at one- and three-second intervals along the length of the garment to direct lymph and extravascular fluids proximally towards the axilla or other functional draining basins within the trunk (Adams, 2010).

In April 2012, the U.S. Food and Drug Administration approved Flexitouch pneumatic compression garment systems for treatment of several clinical indications, including limb lymphedema (U.S. Food and Drug Administration, 2012). In August 2016, the manufacturer received approval to use the Flexitouch garments designed for the head and neck area specifically to treat head and neck lymphedema (U.S. Food and Drug Administration, 2016). Approval was based on data collected from a limited market release to more than 80 adults without active cancer at least four weeks after cancer treatment to determine substantial equivalence. A majority of subjects were able to don and doff the device by the second attempt with a favorable benefit-risk profile of the device for use in the head and neck.

Findings

The American Cancer Society guideline for treating lymphedema after treatment for head and neck cancer makes no mention of pneumatic compression devices (Cohen, 2016). The American Head and Neck Society statement on lymphedema treatment lists several options, including compression bandages and clothing, but not pneumatic compression devices such as Flexitouch (Miller, 2016). A National Comprehensive Cancer Network information guide for patients also does not mention pneumatic compression devices among treatments for lymphedema (National Comprehensive Cancer Network, 2020).

A study ($n = 206$) of subjects with head and neck related lymphedema treated daily for at least 30 minutes with Flexitouch for at least four weeks (average 90 days) revealed significant improvement in all symptoms and function items ($P < .00001$), including ability to control lymphedema at home. A total of 71% of participants

reported compliance with daily use, 87% reported they were satisfied or very satisfied, and 90% reported feeling better after treatment. Authors stated that findings support a randomized controlled trial (Gutierrez, 2020).

Persons with lymphedema who had completed treatment for head and neck cancer were randomized into a group using daily advanced pneumatic compression devices (n = 24) or a wait-list group (n = 25). The group using devices had superior perceived ability to control lymphedema ($P = .003$); less visible external swelling (front view $P < .001$, right view $P = .004$, left view $P = .005$); and less reported pain (Ridner, 2021).

A study of phlebolympheidema treatment compared outcomes for Blue Cross members who used Flexitouch (n = 87) and those who used conservative treatment (n = 86) found the Flexitouch group had 59% fewer mean annual hospitalizations (.13 versus .32; $P < .001$). The study did not mention the original diagnosis, and thus did not specify head and neck cancer cases (Lerman, 2019).

A survey of 44 persons with lymphedema after head and neck cancer therapy subsequently treated with Flexitouch found that 25 rated the device very/somewhat comfortable; 36 rated the treatment very/somewhat comfortable; 27 related feeling somewhat or much better with the device; and 41 stated that home use of the device would be very or somewhat likely (Mayrovitz, 2018).

A systematic review of 26 studies (n = 1,018) of treatment efficacy for lymphedema after head and neck cancer treatment found that few studies are randomized, and manual lymph drainage is best studied. No mention is made of pneumatic compression devices such as Flexitouch (Tyker, 2019).

Upgraded types of pneumatic compression devices for head and neck-related lymphedema are in development. A study of 10 patients involved imaging using near-infrared fluorescence lymphatic imaging before and immediately after advanced pneumatic compression device treatment, and after two weeks of home therapy. Reductions in facial composite measurement scores tracked with reductions in backflow and subject-reported improvements (Gutierrez, 2019).

In 2022, we updated the references, deleted several older references, and identified no new relevant literature to add to the policy. No policy changes are warranted.

References

On January 4, 2022, we searched PubMed and the databases of the Cochrane Library, the U.K. National Health Services Centre for Reviews and Dissemination, the Agency for Healthcare Research and Quality, and the Centers for Medicare & Medicaid Services. Search terms were “head and neck neoplasms (MeSH),” “lymphedema (MeSH),” “Flexitouch,” “head and neck cancer,” “lymphedema,” and “pneumatic compression.” We included the best available evidence according to established evidence hierarchies (typically systematic reviews, meta-analyses, and full economic analyses, where available) and professional guidelines based on such evidence and clinical expertise.

Adams KE, Rasmussen JC, Dame C, et al. Direct evidence of lymphatic function improvement after advanced pneumatic compression device treatment of lymphedema. *Biomed Opt Express*. 2010;1(1):114-125. Doi: 10.1364/BOE.1.000114.

Cohen EE, LaMonte SJ, Erb NL, et al. American Cancer Society head and neck cancer survivorship care guideline. *CA Cancer J Clin*. 2016;66(3):203-239. Doi: 10.3322/caac.21343.

Deng J, Murphy BA, Dietrich MS, et al. Impact of secondary lymphedema after head and neck cancer treatment on symptoms, functional status, and quality of life. *Head Neck*, 2013;35(7):1026-1035. Doi: 10.1002/hed.23084.

Gutierrez C, Karni RJ, Naqvi S, et al. Head and neck lymphedema: Treatment response to single and multiple sessions of advanced pneumatic compression therapy. *Otolaryngol Head Neck Surg.* 2019;160(4):622-626. Doi: 10.1177/0194599818823180.

Gutierrez C, Mayrovitz HN, Naqvi SHS, Karni RJ. Longitudinal effects of a novel advanced pneumatic compression device on patient-reported outcomes in the management of cancer-related head and neck lymphedema: A preliminary report. *Head Neck.* 2020;42(8):1791-1799. Doi: 10.1002/hed.26110.

Lerman M, Gaebler JA, Hoy S, et al. Health and economic benefits of advanced pneumatic compression devices in patients with phlebo-lymphedema. *J Vasc Surg.* 2019;69(2):571-580. Doi: 10.1016/j.jvs.2018.04.028.

Mayrovitz HN, Ryan S, Hartman JM. Usability of advanced pneumatic compression to treat cancer-related head and neck lymphedema: A feasibility study. *Head Neck.* 2018;40(1):137-143. Doi: 10.1002/hed.24995.

Miller M. Lymphedema. AHNS Survivorship, Supportive Care & Rehabilitation Service, American Head & Neck Society. https://www.ahns.info/survivorship_intro/topical_review-lymphedema/. Published 2016.

National Cancer Institute. Head and neck cancers. <https://www.cancer.gov/types/head-and-neck/head-neck-fact-sheet>. Last reviewed May 25, 2021. (a)

National Cancer Institute. Surveillance, Epidemiology, and End Results Program. Cancer stat facts: Oral cavity and pharynx cancer. <https://seer.cancer.gov/statfacts/html/oralcav.html>. Published 2021. (b)

National Comprehensive Cancer Network. Survivorship care for cancer-related late and long-term effects. https://www.nccn.org/patients/resources/life_with_cancer/managing_symptoms/lymphedema.aspx. Published 2020.

Ridner SH, Kietrich MS, Deng J, Ettema SL, Murphy B. Advanced pneumatic compression for treatment of lymphedema of the head and neck: A randomized wait-list controlled trial. *Support Care Cancer.* 2021;29(2):795-803. Epub 2020 Jun 2. Doi: 10.1007/s00520-020-05540-8.

Smith BG, Hutcheson KA, Little LG, et al. Lymphedema outcomes in patients with head and neck cancer. *Otolaryngol Head Neck Surg.* 2015;152(2):284-291. Doi: 10.1177/0194599814558402.

Smith BG, Lewin JS. The role of lymphedema management in head and neck cancer. *Curr Opin Otolaryngol Head Neck Surg.* 2010;18(3):153-158. Doi: 10.1097/MOO.0b013e32833aac21.

Tyker A, Franco J, Massa ST, et al. Treatment for lymphedema following head and neck cancer therapy: A systematic review. *Am J Otolaryngol.* 2019;40(5):761-769. Doi: 10.1016/j.amjoto.2019.05.024.

U.S. Food and Drug Administration. Flexitouch® system. 501(k) Premarket notification letter K120972. https://www.accessdata.fda.gov/cdrh_docs/pdf12/K120972.pdf. Published April 13, 2012.

U.S. Food and Drug Administration. Flexitouch® system. 501(k) Premarket notification letter K153311. https://www.accessdata.fda.gov/cdrh_docs/pdf15/K153311.pdf. Published August 12, 2016.

Policy updates

3/2020: initial review date and clinical policy effective date: 5/2020

4/2020: Policy references updated.

3/2021: Policy references updated.

3/2022: Policy references updated.